

A water soluble granule formulation containing 105 g/kg florasulam, 83 g/kg metsulfuron-methyl and 83 g/kg tribenuron-methyl and for weed control in wheat, barley, oats, rye and triticale.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Maximum individual dose: 50 g/ha
 Maximum total dose: 50 g/ha
 Crops: wheat, barley, oats, rye and triticale
 Latest time of application: GS39



The Advantages:

- 3 active ingredients offering a broad spectrum approach to on time and late application programmes
- All significant cereal crops covered with wide application window
- Broad tank mix recommendations
- Large weed spectrum, and large weeds
- Significant value proposition compared to SU+ Galaxy® mixer



The recommendation
Winter Cereals Clean Up
 AllyULTRA® 50g/ha

Spring Barley
 AllyULTRA® 50g/ha + “Fluroxypyr 200g” 0.6

Targeting the principal problematic weeds in spring cereals (see weed specs over-leaf)

Ally® ULTRA;

Optimum weed control as a stand-alone product for cleaning up, over wintered, groundsel, charlocks, vol.beans and cleavers.
 Or as the main herbicide in spring crops, by adding fluroxypyr

1. Development of resistance with a weed species can be avoided or delayed by alternating (or tank mixing) with suitable products having a different mode of action. For control of Chickweed this product must always be applied with an alternative mode of action product at recommended

© All manufacturers trademarks are duly acknowledged

Ally® Ultra – Weed Specs

	COMMON NAME	Ally Ultra 50g	COMMON NAME	Ally Ultra 50g
Label Weeds	Black Bindweed ★★	6 lf	Fumitory, common	MS
	Cleavers ★	S	Greater Burdock	90 cm
	Common Chickweed** ★	> 15 cm*	Greater Plantain	ms
	Fat-hen	> 2 lf	Green Field Speedwell	2 lf
	Field Pansy	6 lf	Groundsel	10 cm
	Knotgrass	> 6 lf	Hedge mustard	S < 10cm
	Pale Persicaria ★	15 cm	Hemlock	yes
	Poppy Common	known resistance	Hemp-nettle	4 tl
	Red Dead-nettle ★	15 cm	Henbit Dead-nettle ★	6 lf
	Redshank	15 cm	Hogweed	F > 30 cm
	Scented Mayweed	S - upto Flower bud visible	Ivy-leaved Speedwell	2 lf 5cm
	Scentsless Mayweed	S - upto Flower bud visible	Mouse-ear Chickweed	20 cm
	Shepard's Purse	> 15 cm	Nipplewort	6lf
	Speedwell, common field	ms 8etl	Parsley Piert	15 cm
	Speedwell, ivy leaf	ms cotyledon	Perennial Sow-thistle	R 4lf/8cm
Volunteer Rape	> 15 cm	Pigweed	10 cm	
Background Weeds (non label)	Black Nightshade ★	5cm	Pineappleweed	11 cm
	Broad-leaved Dock	20-50 cm	Rosebay Willowherb	MS
	Bugloss	2 lf	Runch	> 15 cm
	Charlock	before flower bud visible	Scarlet Pimpernel	15 cm
	Coltsfoot	2lf	Sheep's Sorrel	MS
	Common Comfrey	MS	Shepherd's Needle	> 15 cm
	Common Field speedwell	> 6 lf/20 cm	Silverweed	10 cm
	Common Fumitory	MR-R	Slender Speedwell	2 lf
	Common Hemp-nettle	> 15 cm	Small Nettle	6 lf
	Common Orache	6 lf	Smooth Hawk's-beard	F 12 cm
	Common Poppy ★	> 6 lf	Sow-thistle, annual	2-4 leaf
	Common Yetch	10 cm	Sow-thistle, smooth	6lf
	Corn Buttercup	ms	Speedwell, wall	2 lf
	Corn Chamomile	F 20 cm	Spring Beauty	15 cm
	Corn Cockle	4 lf	Square-stalked Willowherb	MS
	Corn Marigold		Stinging Nettle	15 cm
	Corn Mint	4 lf	Stinking Chamomile	S - upto Flower bud visible
	Corn Spurrey	15 cm	Sun Spurge	> 2 lf
	Cornflower	2 lf	Tar Weed	6 lf
	Cow Parsley	yes	Thale cress	6tl
	Creeping Buttercup	2 lf	Venus's Looking glass	15 cm
	Creeping Thistle	F yes	Volunteer Beans	10cm
	Curled Dock ★	20-50 cm	Volunteer Lucerne	sd
	Dandelion	F > 4 lf	Volunteer Potatoes	R 5-10 cm (needs Fluroxypyr for top growth)
	Docks (from seed) ★	S	Volunteer Sugar Beet	15cm
	Dovesfoot Cranesbill	6 lf	White Campion	4 lf
	Field Forget-me-not	6 lf	Wild carrot	> 15 cm
	Field Horsetail	R 8-10 cm	Wild Garlic/Ramsons	yes
	Field Pennygrass	yes	Wild Mignonette	2 lf
	Flixweed	yes	Wild Onion	yes
	Fool Parsley	6 lf	Wild Radish (Runch)	> 15 cm

Now add in the tank mix partner “Fluroxypyr” and control of the weeds highlighted with ★ are significantly enhanced (including, Nightshade, red dead nettle, henbit dead nettle, vol. linseed

Most marigolds, we assume to be resistant, and require the inclusion of clopyralid

★ Increasing numbers of common poppy are presenting as resistant, an alternative mode of action to ALS is required

1. Development of resistance with a weed species can be avoided or delayed by alternating (or tank mixing) with suitable products having a different mode of action. For control of Chickweed this product must always be applied with an alternative mode of action product at recommended

© All manufacturers trademarks are duly acknowledged