



## **CURZATE<sup>®</sup> M WG fungicide**

**PCS NO 04723**

A water dispersible granule containing 4.5% w/w cymoxanil and 68% w/w mancozeb for the control of potato late blight in potatoes

### **SAFETY PRECAUTIONS**

#### **ENVIRONMENTAL PROTECTION**

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

#### **RISK AND SAFETY INFORMATION**

### **Curzate<sup>®</sup> M WG**

#### **Fungicide**

Containing 4.5% w/w cymoxanil and 68% w/w mancozeb

Contains cymoxanil and mancozeb. May produce an allergic reaction

#### **WARNING**



**May cause an allergic skin reaction.**

**Suspected of damaging the unborn child.**

**Very toxic to aquatic life with long lasting effects.**

Avoid breathing vapours/spray.

Wear protective gloves / protective clothing.

If medical advice is needed, have product container or label at hand

Contaminated work clothing should not be allowed out of the workplace

**IF ON SKIN:** Wash with plenty of soap and water.

**IF exposed or concerned:** Get medical advice/attention.

**If skin irritation or rash occurs:** Get medical advice/attention.

Take off contaminated clothing and wash it before re-use

Keep out of reach of children

Store locked up

Collect spillage.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste

**To avoid risks to human health and the environment, comply with the instructions for use**

**PCS No. 04723**

Contents: XX Kg e

Batch No: XXXX  
Manufactured in EU

**Du Pont (U.K.) Limited**  
**Crop Protection Products**  
**4th Floor**  
**Kings Court**  
**London Road**  
**Stevenage**  
**Herts SG1 2NG**

**Tel: 00 44 1438 734450 or enquiry.agproducts@dupont.com**  
**National Poisons Information Centre (24hr): 01 837 9964 or 01 809 2166**  
**DuPont Emergency Contact (24hr): (01) 901 4670**

® Registered trademark of E.I. du Pont de Nemours and Company  
FOR PROFESSIONAL USE ONLY

## DIRECTIONS FOR USE

### RESTRICTIONS

- Allow a minimum of 7 days between applications
- Do not allow packs to become wet during storage
- Do not keep opened packs from one season to another
- Keep away from fire or sparks

### DISEASE CONTROL

CURZATE M WG combines the local systemic action of cymoxanil and the protectant activity of mancozeb for the control of potato late blight (*Phytophthora infestans*) of potatoes

#### Disease Resistance

When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. It is therefore advised that wherever possible spray programmes should include fungicides with differing modes of action.

### CROPS

CURZATE M WG can be used on all varieties of early and maincrop potatoes including seed crops. On crops grown for processing consult your processor before use.

### TIMING AND WEATHER

Timing of spray applications should take into account the normal prevalence of the disease in the area and conditions in the particular season. The first spray should be applied as soon as there is a risk of blight infection or an official warning has been issued. In the absence of a warning, spraying should begin just before the crop meets along the rows.

Applications should be repeated at 10-14 day intervals depending on the prevalence of the disease and favourable conditions for infection. A 7 day spray interval may be used under exceptional circumstances e.g. blight risk is already high and rain is forecast at the time of the next 10 day spray

is due. The longer interval is recommended under dry conditions. Irrigation of crops increases the risk of blight and crops should always be sprayed with CURZATE M WG as soon as possible after irrigation subject to the minimum spray interval. High-risk situations include conditions of rapid growth, irrigated crops, areas of intensive potato growing, particularly blight-sensitive varieties, when further warnings are issued, or when active blight is in the area.

## DOSE

Apply CURZATE M WG at 2.3 kg/ha

## VOLUME AND APPLICATION

Apply CURZATE M WG in not less than 200 litres of water per hectare using a MEDIUM quality spray (as defined by BCPC)

## MIXING

CURZATE M WG mixes easily with water, but the following mixing procedure should be followed: Quarter fill the spray tank with water and start the agitation. Slowly add the required quantity of CURZATE M WG directly to the tank. Continue agitation while topping up the tank and while spraying.

## WARNING

KEEP AWAY FROM FIRE OR SPARKS.

Do not allow CURZATE M WG to become wet during storage or keep opened packs from one season to another as this may reduce the effectiveness of the fungicide and may create vapours that may be flammable.

## NOTICE TO BUYER

All goods supplied by us are of high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

### FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Max single dose (L/ha)	Max no of applications	Max total dose (L/ha)	Latest time of application
Potato (seed and ware)	2.3kg/ha	6	13.8kg/ha	14 days before harvest

**Method of application:** Tractor mounted / trailed sprayer

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**