

IRL

# SUPER RAPEEZE MSO

FOR USE ONLY AS AN AGRICULTURAL HORTICULTURAL INDUSTRIAL FORESTRY ADJUVANT OIL		
Crops/situations:	Pesticides	Maximum concentration
All non edible crops	All approved pesticides	1.78% spray solution
All edible crops	All approved pesticides at half or less than half the approved pesticide rate	1.78% spray solution
For maximum number of treatments and latest time of application, follow the statutory conditions of use of pesticide. Read all other safety precautions and directions before use		

**ACTION: INTRACROP SUPER RAPEEZE MSO** is an adjuvant containing methylated rapeseed oil, which readily emulsifies in water. It has low viscosity and is easy to mix giving a stable and homogeneous emulsion with a wide range of pesticides.

**SUPER RAPEEZE MSO** improves the wetting, spreading and sticking properties of crop protection products on leaf surfaces.

**DIRECTIONS FOR USE:**

**RATES OF USE - SUPER RAPEEZE MSO** is recommended at a rate of use of up to 2.0 litres product per hectare.

**VOLUME:** 112 - 300 litres of water per hectare. For water volumes below 200 l/ha, a concentration of 1% in the final spray solution is ideal. Do not exceed a concentration of 1.78%. Always maintain continuous agitation during spraying.

**MIXING:** Follow any specific mixing instructions for the partner pesticide.

Add the **SUPER RAPEEZE MSO** to the spray mix in the sprayer tank **BEFORE** filling up the sprayer with water.

Continue agitation whilst filling the sprayer tank with water.



**INTRACROP®**

**SAFETY PRECAUTIONS:**

**Operator Protection:** Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment. \*Wear suitable protective gloves and eye/face protection (faceshield) when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

Do not get in eyes, on skin, or on clothing. Do not eat, drink or smoke when using this product.

Wash hands thoroughly after handling. IF exposed or if you feel unwell: Call a POISON CENTRE or doctor.

**Environmental Protection:** Do not use with any pesticide, which is to be applied in or near water.

**Storage and Disposal:** Keep locked and out of reach of children. Keep away from food, drink and animal feeding stuffs.

Wash out container thoroughly and dispose of safely. Do not re-use the container for any purpose.

<b>SUPER RAPEEZE MSO</b> may be used with all approved crop protection products at recommended rates of use up to the growth stages below. Once the crops have grown past the stages, application rate must reduce to half the approved pesticide rate	
Crop	Growth stages before which full rates can be used
Apples / pears	up to and including fruit 5 – 10mm
Cherry / plum	up to and including first fruit set
Almond, apricot and peach	up to and including first fruit set and peach
Table grapes and wine grapes	up to and including first fruit set
Strawberry	up to and including first fruit development
Raspberry	up to and including first fruit set
Blackcurrant	up to and including first fruit visible green fruitlet
Cucumber	up to and including first fruit set
Tomato	up to and including first fruit set on truss
Combining and vining peas	up to and including flat pod set (non edible pods only)
Edible podded peas	up to and including pod set
Oil-seed rape	up to and including 10% potential pods
Linseed	up to and including 10% capsules formed
Sugar beet	up to and including 6 fully expanded leaves
Hops	up to and including cone set
Cereals	up to and including GS52 (¼ inflorescence)
Brussels sprout	up to and including lateral buds begin to develop
Maize	up to and including tip of tassel visible
Potatoes	up to and including tuber initiation
Beans	up to and including first pod set
Cauliflower	up to and including cauliflower heads begin to form; width of growing tip up to 1cm
Lettuce	Varieties forming heads: up to and inc. heads begin to form Varieties not forming heads: up to and including leaf rosette reached 30% of diameter typical for the variety
When crops have passed the growth stages above, only use <b>SUPER RAPEEZE MSO</b> with HALF or LESS THAN HALF of the recommended rate of use of the pesticide.	

**CONDITIONS OF SUPPLY :**

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or our re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

SRP/15/01/IRL/BJ