



Directions for use: INCITE* may be

applied to all crops as a foliar spray at

a rate of 0.5 - 2 litres per hectare. Úse

sufficient water volume to ensure good

coverage - typically 200 - 400 litres per

INCITE* is intended as a supplement to

a regular fertiliser programme and will

crop recommendations. For all round

protection and weather proofing add

Nu Film P at a rate of 120ml per hectare.

Consult INTRACROP for specific

not on its own provide full crop nutrition.

hectare.

INCITE/L/010/02/IE

WHERE IT IS INTENDED TO MAKE **ONLY A SINGLE APPLICATION USE** THE HIGH RATE

Mixing: Shake the bottle well before use. Half fill the sprayer with water, add the required quantity of INCITE* and fill with the remainder of the water. Place the INCITE* into the tank before adding any other chemicals.

Compatibility: INCITE* is compatible with many pesticides, fertilisers and surfactants. Before applying tank mixes to the crop, first check safely on a small area. Do not apply INCITE* together with alkaline materials, crop oil adjuvants or penetrating type surfactants.

All goods supplied by us are of high grade and we believe them to be suitable for the purpose recommended. Products are supplied solely for the use and applications recommended subject to strict adherence to the label conditions of storage and use. Other uses. applications and deviations from label recommendations are solely at the user's own risk since we cannot exercise control over storage or seasonal factors affecting weather conditions, growth stage, crop conditions and the composition of tank mixtures. No liability or warranty is expressed or implied concerning the safety of tank mixtures or their biological effects on crop safety.

APPLICATION RATE CROP **TIMING** Litres per Hectare Beans 4 nodes to early pod formation (Use higher dose at pod formation) 1 - 2 Carrots When storage root growth commences 1 - 2 **Brussel Sprouts** Button stage 1 - 2 Cereals 21/2 leaves to Zadox 32 (2nd node) 0.5 - 12½ leaf to early tillering 1 - 2 Grass As soon as new growth commences after cutting 1 - 2 Oilseed Rape 5 true leaves to early stem elongation 1 Onions/Leeks 2 3 to 6 true leaves 1 2 fully expanded leaves to 3 pairs of fully expanded leaves Peas 1 - 2 Flower bud formation (best timing) to early pod formation 7 - 10 days before flowering commences or when 2 - 4 tubers are between 0.6 and 2.4 cm in diameter **Potatoes** 2 Further application 3 - 5 weeks later 1 - 2 3 to 5 leaves Sugar Beet When storage root development becomes marked 2 (from about 8 leaves) Maize When crop is at the 4 - 8 leaf stage 1 - 2 Swedes/Turnips 6 to 10 leaf stage 1 - 2 Vegetable/ 6 to 8 true leaves 1 - 2 Brassicas

Statutory Fertiliser Declaration All values given as W/w %, for W/v % or g/L please refer to product literature or contact the supplier INCITE® is a 0 - 28.5 - 20.5 PK Fertiliser Solution Water soluble Phosphorous Pentoxide (P2O5) 28.5 % (12.5 % P) Water soluble Potassium Oxide (K2O) 20.5 % (17.0 % K)

Health & Safety Information H318: Causes serious eve damage.

P264: Wash hands, forearms and face thoroughly after handling.

P280: Wear protective gloves/ protective clothing/eye protection/face protection.

P305 +P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

P309 +P311: IF exposed or if you feel unwell: Call a POISÓN CENTRE or doctor/physician.

P502: Refer to manufacturer or supplier for information on recovery or recycling.



TERRACHEM LTD

Tel: +353 87 254 7534 Email: info@terrachem.ie www.terrachem.ie



INTRACROP* Park Chambers, 10 Hereford Rd, Abergavenny, Monmouthshire, NP7 5PR, UK | Email: admin@intracrop.co.uk Tel: +44 (0)1926 634 801 | www.intracrop.co.uk | **9** @intracrop 24Hr Emergency ONLY Tel: +44 (0)870 190 6777